

KARMEN Limits on $\nu_e \rightarrow \nu_\tau$ Oscillations in 2- ν and 3- ν Mixing Schemes

B. Armbruster^a, I. Blair^b, B.A. Bodmann^c, N.E. Booth^d, G. Drexlin^a, V. Eberhard^a, J.A. Edgington^b, C. Eichner^e,
K. Eitel^a, E. Finckh^c, H. Gemmeke^a, J. Hößl^c, T. Jannakos^a, P. Jünger^c, M. Kleifges^a, J. Kleinfeller^a,
W. Kretschmer^c, R. Maschuw^e, C. Oehler^a, P. Plischke^a, J. Rapp^a, C. Ruf^e, M. Steidl^a, O. Stumm^c, J. Wolf^{a*},
B. Zeitnitz^a

^a *Institut für Kernphysik I, Forschungszentrum Karlsruhe, Institut für experimentelle Kernphysik, Universität Karlsruhe, Postfach 3640, D-76021 Karlsruhe, Germany*

^b *Physics Department, Queen Mary and Westfield College, Mile End Road, London E1 4NS, United Kingdom*

^c *Physikalisches Institut, Universität Erlangen-Nürnberg, Erwin Rommel Strasse 1, D-91058 Erlangen, Germany*

^d *Department of Physics, University of Oxford, Keble Road, Oxford OX1 3RH, United Kingdom*

^e *Institut für Strahlen- und Kernphysik, Universität Bonn, Nußallee 14-16, D-53115 Bonn, Germany*

The 56 tonne high resolution liquid scintillation calorimeter KARMEN at the beam stop neutrino source ISIS has been used to search for neutrino oscillations in the disappearance channel $\nu_e \rightarrow \nu_x$. The ν_e emitted in μ^+ decay at rest are detected with spectroscopic quality via the exclusive charged current reaction $^{12}\text{C}(\nu_e, e^-)^{12}\text{N}_{\text{g.s.}}$ almost free of background. Analysis of the spectral shape of e^- from the ν_e -induced reaction as well as a measurement of the absolute ν_e flux allow to investigate oscillations of the type $\nu_e \rightarrow \nu_\tau$ and $\nu_e \rightarrow \nu_\mu$. The flux independent ratio $R_{CC/NC}$ of charged current events $^{12}\text{C}(\nu_e, e^-)^{12}\text{N}_{\text{g.s.}}$ to neutral current events $^{12}\text{C}(\nu, \nu')^{12}\text{C}^*$ provides additional information in the oscillation channel $\nu_e \rightarrow \nu_x$. All three analysis methods show no evidence for oscillations. For the $\nu_e \rightarrow \nu_\tau$ channel 90% CL limits of $\sin^2(2\theta) < 0.338$ for $\Delta m^2 \geq 100 \text{ eV}^2/\text{c}^4$ and $\Delta m^2 < 0.77 \text{ eV}^2/\text{c}^4$ for maximal mixing in a simple 2 flavor oscillation formalism are derived. A complete 3 flavor analysis of the experimental data from five years of measurement with respect to $\nu_e \leftrightarrow \nu_\tau$ and $\nu_e \leftrightarrow \nu_\mu$ mixing is presented.

I. INTRODUCTION

One of the most interesting issues in present particle physics is to clarify the neutrino mass problem. In the minimal standard model of electroweak theory neutrinos are considered to be massless. But there is no compelling theoretical reason behind this assumption. On the other hand, in most extensions of the standard model massive neutrinos are allowed. A very sensitive way of probing small neutrino masses and the mixing between different neutrino flavors is provided by neutrino oscillations. Although there are some experimental results pointing to neutrino oscillations, it is still difficult to deduce a consistent and reliable set of neutrino masses and mixings [1–3].

Up to now, experimental results have been persistently interpreted in terms of neutrino oscillations using only a 2- ν mixing scheme with a single mixing angle Θ . However, to perform a complete and consistent analysis of experimental data on neutrino oscillation search, a 3 flavor formalism of neutrinos should be adopted. In this framework, all experiments searching for different flavor oscillations (e.g. $\nu_\mu \rightarrow \nu_e$, $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$, $\nu_\mu \rightarrow \nu_\tau$ in appearance mode or $\nu_e \rightarrow \nu_x$, $\bar{\nu}_e \rightarrow \bar{\nu}_x$, $\nu_\mu \rightarrow \nu_x$ in disappearance mode) are combined to extract global information on the oscillation parameters, the mixing angles as well as the mass differences $\Delta m^2_{ij} = |m_i^2 - m_j^2|$, $i, j = 1, \dots, 3$. Searches for oscillations in the disappearance mode, although less sensitive to small mixing angles, are of great importance to restrict the allowed parameter space.

In many of the 3 flavor descriptions of neutrino oscillations a so-called 'one-mass-scale dominance' $\delta m^2 \equiv \Delta m^2_{12} \ll \Delta m^2_{13}, \Delta m^2_{23} \approx \Delta m^2_{23} \equiv \Delta m^2$ has been adopted [3–9]. Possible mixing to sterile neutrinos as suggested by [10–12] is ignored whereas CP conservation is assumed, as we shall do in the following. The flavor eigenstates ν_α are then described by superpositions of the mass eigenstates ν_i

$$\nu_\alpha = \sum_{i=1}^3 U_{\alpha i} \nu_i \quad (\alpha = e, \mu, \tau) \quad (1)$$

with the mixing matrix elements $U_{\alpha i}$ following the typical notation [13] of the CKM-like mixing matrix described by two mixing angles Φ, Ψ . The oscillation probability $P_{\alpha\beta}$ for transition $\nu_\alpha \rightarrow \nu_\beta$ in the KARMEN experiment with its 'short baseline' experimental configuration ($1/\Delta m^2 \approx L/E \ll 1/\delta m^2$) is then

$$P_{\alpha\beta} = A \cdot \sin^2(1.27 \frac{\Delta m^2 L}{E}) \quad (2)$$

with Δm^2 in eV^2/c^4 , the neutrino path length L in meters and the neutrino energy E in MeV. In a simple 2 flavor description $A = \sin^2(2\Theta)$ with Θ being the mixing angle between the two ν states, while in a 3 flavor analysis $A = 4U_{\alpha 3}^2 U_{\beta 3}^2$ (appearance mode) or $A = 4U_{\alpha 3}^2(1 - U_{\alpha 3}^2)$ (disappearance mode) with $U_{e3}^2 = \sin^2\Phi$, $U_{\mu 3}^2 = \cos^2\Phi \sin^2\Psi$, $U_{\tau 3}^2 = \cos^2\Phi \cos^2\Psi$ [3]. For an experimental oscillation search it is important to note that the second factor in (2) describing the spatial evolution of oscillations is unchanged in the 2 flavor analysis. In the deduction of limits on A and Δm^2 we shall start with $A = \sin^2(2\Theta)$ having in mind that the oscillation amplitude A has different definitions depending on the actual theoretical model as well as the experimental situation (appearance $\leftrightarrow$ disappearance search). In section IV and V we shall then analyze our results in the 3 flavor scheme.

The **K**arlsruhe **R**utherford **M**edium **E**nergy **N**eutrino experiment KARMEN has set stringent upper limits in the appearance channels $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ ($\sin^2(2\Theta) < 8.5 \cdot 10^{-3}$ for $\Delta m^2 \geq 100 \text{eV}^2/c^4$) and $\nu_\mu \rightarrow \nu_e$ ($\sin^2(2\Theta) < 4.0 \cdot 10^{-2}$ for $\Delta m^2 \geq 100 \text{eV}^2/c^4$) [14]. KARMEN also searches for $\nu_e \rightarrow \nu_\tau$ and $\nu_e \rightarrow \nu_\mu$ through disappearance of ν_e . Here, the neutrinos ν_e emerge from μ^+ decay at rest (DAR) and are detected at a mean distance of 17.7 m from the source. This search is a complementary addition to the appearance channels. Compared to other disappearance experiments this search has very small, systematic uncertainties as well as an excellent signal to background ratio for the detection of ν_e -induced events. Limits on $\bar{\nu}_e$ disappearance were obtained by reactor experiments e.g. at the nuclear power plant of Bugey [15] with $\bar{\nu}_e$ from β -decays and the detector at 15, 40 or 95 m distance. Other limits on $\nu_e \rightarrow \nu_x$ were deduced from two ^{51}Cr source experiments in the GALLEX detector [16,17] emitting ν_e with discrete energies $E_\nu = 0.746 \text{ MeV}$ (81%), 0.751 MeV (9%), 0.426 MeV (9%) and 0.431 MeV (1%). Experiments with neutrinos in the energy range of several GeV allow the search for appearance $\nu_e, \nu_\mu \rightarrow \nu_\tau$. In contrast to such experiments, like FNAL E531 [18] with a 410 meter long decay path, in KARMEN the parameters L and E of ν -induced events can be measured with high accuracy on an event-by-event basis.

The outline of the paper is as follows: We first present the experimental setup and the ν_e detection and identification (section II). Then we investigate the absolute number of identified CC reactions $^{12}\text{C}(\nu_e, e^-)^{12}\text{N}_{\text{g.s.}}$ (section III A) comparing the measured cross section with theoretical predictions. In a second evaluation method (section III B) we normalize the number of CC sequences to the number of neutral current (NC) reactions $^{12}\text{C}(\nu, \nu')^{12}\text{C}^*$ (15.1 MeV). These NC reactions are not sensitive to the ν flavor. By comparing the number of CC events with NC events measured simultaneously, systematic uncertainties in the calculation of the absolute ν flux mostly cancel. A lower ratio of events $R_{\text{exp}} = \text{CC}/\text{NC}$ than expected would also point to $\nu_e \rightarrow \nu_x$. A complementary analysis is given in section III C, where we search for distortions of the energy and spatial distributions of electrons from $^{12}\text{C}(\nu_e, e^-)^{12}\text{N}_{\text{g.s.}}$, such as would be due to $\nu_e \rightarrow \nu_x$. In section IV the results of these analyses are combined and compared with other experimental limits for $\bar{\nu}_e \rightarrow \bar{\nu}_x$, $\nu_e \rightarrow \nu_\tau$ and $\nu_\mu \rightarrow \nu_e$ ($\bar{\nu}_\mu \rightarrow \bar{\nu}_e$) in a 2 flavor as well as a 3 flavor mixing scheme.

II. KARMEN EXPERIMENT

A. Experimental Setup

The KARMEN experiment is performed at the neutron spallation facility ISIS of the Rutherford Appleton Laboratory. The rapid cycling synchrotron provides 800 MeV protons with an average current of $200 \mu\text{A}$. These are stopped in a beam-stop Ta-D₂O-target producing neutrons and pions. 99% of all charged pions are stopped within the target [19] leading to π^+ decay at rest whereas negative pions are absorbed in the heavy target nuclei. Neutrinos therefore emerge from the consecutive decay sequence $\pi^+ \rightarrow \mu^+ + \nu_\mu$ and $\mu^+ \rightarrow e^+ + \nu_e + \bar{\nu}_\mu$ in equal intensity per flavor. The intrinsic contamination $\bar{\nu}_e/\nu_e$ from μ^- decay in the beam-stop-target is only 4.7×10^{-4} [19]. The neutrino energy spectra are well defined due to the decay at rest kinematics: The ν_μ from π^+ -decay are monoenergetic ($E_\nu = 29.8 \text{ MeV}$), the continuous energy distributions of ν_e and $\bar{\nu}_\mu$ up to 52.8 MeV can be calculated using V-A theory (Fig. 1a) where the characteristic Michel shape of ν_e from μ^+ DAR is given by the expression [20]

$$N(\epsilon)d\epsilon = 4\epsilon^2 [3(1-\epsilon) + \frac{2}{3}\rho(4\epsilon-3)]d\epsilon \quad \text{with } \epsilon = \frac{E_\nu}{E_{\text{max}}} \text{ and } \rho = 0 \text{ for } \nu_e \quad (3)$$

The neutrino production time follows the two parabolic proton pulses of 100 ns base width and a separation of the maxima of 324 ns, produced with a repetition frequency of 50 Hz. The different lifetimes of pions ($\tau = 26 \text{ ns}$) and muons ($\tau = 2.2 \mu\text{s}$) together with this unique time structure allow a clear separation in time of the ν_μ -burst from the following ν_e and $\bar{\nu}_\mu$ (Fig. 1b) which show the characteristic exponential distribution of the muon lifetime $\tau = 2.2 \mu\text{s}$. Furthermore the accelerator's extremely small duty cycle suppresses effectively any beam-uncorrelated background by four to five orders of magnitude depending on the time interval selected for analysis.

The neutrinos are detected in a high resolution 56 t liquid scintillation calorimeter segmented into 512 central modules with a cross section of $0.18 \times 0.18 \text{ m}^2$ and a length of 3.53 m each [21]. A massive blockhouse of 7000 t of

steel in combination with a system of two layers of active veto counters provides shielding against beam correlated spallation neutron background and suppression of the hadronic component of cosmic radiation, as well as highly efficient identification of cosmic muons and their interactions. The central scintillation calorimeter and the inner veto counters are segmented by double acrylic walls with an air gap providing efficient light transport via total internal reflection of the scintillation light at the module walls. The event position within an individual module is determined with a position resolution of ± 6 cm for typical energies of 30 MeV [22] by the time difference of the PM signals at each end of this module. Due to the optimized optical properties of the organic liquid scintillator [23] and an active volume of 96% for the calorimeter, an energy resolution of $\sigma_E = 11.5\%/\sqrt{E[\text{MeV}]}$ is achieved. The KARMEN electronics is synchronized to the ISIS proton pulses to an accuracy of better than ± 2 ns, so that the time structure of the neutrinos can be exploited in full detail.

B. Detection and Identification of ν_e

Electron neutrinos ν_e from μ^+ decay at rest (DAR) are detected via the exclusive charged current (CC) reaction $^{12}\text{C}(\nu_e, e^-)^{12}\text{N}_{\text{g.s.}}$ which leads to a delayed coincidence signature. The ground state of ^{12}N decays at its production point with a lifetime of $\tau = 15.9$ ms: $^{12}\text{N}_{\text{g.s.}} \rightarrow ^{12}\text{C} + e^+ + \nu_e$. The detection signature consists of a prompt electron within a few μs after beam-on-target ($0.6 \leq t_{\text{pr}} \leq 9.6 \mu\text{s}$) followed by a spatially correlated positron (e^-/e^+ sequence with $0.5 \leq t_{\text{diff}} \leq 36$ ms). To reduce cosmic background, no event is accepted within a software deadtime of $20 \mu\text{s}$ after any signal in the detector system. Remaining sequences induced by cosmic background are studied with high statistical accuracy by demanding the prompt event to emerge in a $200 \mu\text{s}$ long time interval before beam-on-target when no neutrinos are produced at ISIS. This CC reaction has been previously studied in detail [24], with respect to weak nuclear form factors [25] and to the CC helicity structure of muon decay [26].

The data set used for this $\nu_e \rightarrow \nu_x$ analysis is taken from 1990–1995 corresponding to 9122 C protons on the ISIS beam stop target, or 2.51×10^{21} μ^+ decays at rest in the target. Figure 2 shows the signatures of 513 events surviving all cuts. These events contain 499.7 ± 22.7 ν -induced e^-/e^+ sequences and 13.3 ± 0.8 cosmic induced background sequences. The energy spectra of the prompt e^- and the delayed e^+ follow the Monte Carlo (MC) expectations from μ^+ DAR and the ^{12}N β -decay. The time distribution of the prompt event reflects the muon lifetime, the time difference t_{diff} shows the ^{12}N lifetime folded with the detection efficiency.

The stringent signatures of the $^{12}\text{C}(\nu_e, e^-)^{12}\text{N}_{\text{g.s.}}$ reaction lead to an excellent signal-to-background ratio of 37:1 which allows identification of ν_e almost free of background. Accordingly, the search for ν_e disappearance has very small systematic errors and the sensitivity in this oscillation channel is not limited by background events.

III. OSCILLATION SEARCHES

Flavor oscillations of the type $\nu_e \rightarrow \nu_\mu$ or $\nu_e \rightarrow \nu_\tau$ will result in a smaller number of detected $^{12}\text{C}(\nu_e, e^-)^{12}\text{N}_{\text{g.s.}}$ sequences than expected because the analogous charged current reactions $^{12}\text{C}(\nu_\mu, \mu^-)^{12}\text{N}_{\text{g.s.}}$ and $^{12}\text{C}(\nu_\tau, \tau^-)^{12}\text{N}_{\text{g.s.}}$ cannot be induced due to the small neutrino energies ($E_\nu \leq 52.8$ MeV). In addition to the comparison of the observed ν_e rate with expectation, the effects of ν_e disappearance can be investigated by a detailed study of the remaining ν_e -induced events. First, the characteristic energy dependence of flavor oscillations (2) will lead to a distortion of the electron energy distribution. Secondly, the $1/r^2$ dependence of the ν_e flux from μ^+ DAR in the beam stop target will be distorted. The specific features of neutrino production at ISIS and detection by KARMEN require the two oscillation channels $\nu_e \rightarrow \nu_\mu$ and $\nu_e \rightarrow \nu_\tau$ to be discussed separately.

We first consider transitions $\nu_e \rightarrow \nu_\tau$. As described above, ν_τ cannot be detected via CC reactions, so oscillations $\nu_e \rightarrow \nu_\tau$ directly lead to a reduction of CC sequences and consequently of the measured flux averaged cross section $\langle \sigma_{\text{CC}} \rangle_{\text{exp}}$.

The second channel, $\nu_e \leftrightarrow \nu_\mu$ mixing, is more complex due to the simultaneous presence of $\bar{\nu}_\mu$ in the neutrino beam from μ^+ decay. Assuming CPT invariance, there would be a second source for CC reactions in the KARMEN detector, namely $\bar{\nu}_e$ from $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ leading to $^{12}\text{C}(\bar{\nu}_e, e^+)^{12}\text{B}$ since $\bar{\nu}_\mu$ from μ^+ DAR at ISIS are produced with the same intensity and at the same time as ν_e . These would mostly compensate the reduction through $\nu_e \rightarrow \nu_\mu$ of the measured cross section due to the following reasons:

Because ^{12}B belongs to the isobar triplet ^{12}B - ^{12}C - ^{12}N , the Q-value for $^{12}\text{C}(\bar{\nu}_e, e^+)^{12}\text{B}$ ($Q = -13.4$ MeV) as well as the ^{12}B lifetime ($\tau = 29.14$ ms) and the endpoint energy of $^{12}\text{B}_{\text{g.s.}} \rightarrow ^{12}\text{C} + e^- + \bar{\nu}_e$ ($E_0 = 13.4$ MeV) are very similar to the corresponding quantities for $^{12}\text{C}(\nu_e, e^-)^{12}\text{N}_{\text{g.s.}}$ with the subsequent ^{12}N β -decay (see Fig. 3). Moreover, the cross sections are similar with $\langle \sigma \rangle = 9.0 \cdot 10^{-42} \text{ cm}^2$ [27] induced by $\bar{\nu}_e$ from $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ with large Δm^2

(see Fig. 4). So, as one CC reaction disappears, the other takes its place.¹ Figure 5 demonstrates this argument in detail showing the relative detection efficiencies $\epsilon(\nu)$ and $\epsilon(\bar{\nu})$ as a function of the oscillation parameter Δm^2 . Taking into account the different energy dependent cross sections for both CC reactions, the detection efficiency $\epsilon(\nu, \Delta m^2)$ within the KARMEN detector for $\nu_e \rightarrow \nu_x$ disappearance into ν_τ or ν_μ is only slightly higher than for $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ appearance $\epsilon(\bar{\nu}, \Delta m^2)$ which also incorporates the different detection efficiency for the sequential e^+/e^- due to the different lifetimes and Q-values of $^{12}\text{N}/^{12}\text{B}$. The different ν energy spectra and energy dependences of the cross sections are responsible for a small shift in the position of minima and maxima of the efficiencies for $\nu_e \rightarrow \nu_x$ and $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$. Therefore, at $\Delta m^2 \approx 5 \text{ eV}^2/\text{c}^4$ a small net increase of CC reactions is expected assuming mixing between ν_e and ν_μ .

Following these considerations the performed search for $\nu_e \rightarrow \nu_x$ must actually be interpreted in terms of $\nu_e \leftrightarrow \nu_\mu$ and $\nu_e \leftrightarrow \nu_\tau$ mixing where the $\nu_e \rightarrow \nu_\tau$ search is much more sensitive (in terms of the absolute number of CC events; see section III A) than $\nu_e \rightarrow \nu_\mu$ due to the almost complete cancellation by the mirror $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$. An analysis of the spectral shape of CC sequences, however, is sensitive to both mixings, $\nu_e \leftrightarrow \nu_\tau$ and $\nu_e \leftrightarrow \nu_\mu$, as will be shown in section III C. This shape analysis is performed in a complete 3 flavor scheme where the amplitudes for the mixings $\nu_e \leftrightarrow \nu_\tau$ and $\nu_e \leftrightarrow \nu_\mu$ are completely free and independent parameters.

A. Absolute Cross Section

The comparison of the number of ν_e -induced CC reactions with the expectation from theoretical calculations is done on the basis of the measured flux averaged cross section $\langle \sigma_{\text{CC}} \rangle_{\text{exp}}$ for ν_e from μ^+ DAR. Taking into account the overall detection efficiency of $\epsilon_{\text{CC}} = 0.328$ the 499.7 ± 22.7 ν_e -induced events (see Fig. 2) correspond to a flux averaged cross section of

$$\langle \sigma_{\text{CC}} \rangle_{\text{exp}} = (9.4 \pm 0.4_{(\text{stat.})} \pm 0.8_{(\text{syst.})}) \times 10^{-42} \text{ cm}^2 \quad (4)$$

where the systematic error is almost entirely due to the uncertainty in the calculation of the absolute ν flux from ISIS [19]. The measured cross section is in excellent agreement with theoretical calculations based on different models to describe the ^{12}C nucleus (see table I), i.e. there is no indication of ν_e disappearance. For quantitative calculations we use the mean theoretical value of $\langle \sigma_{\text{CC}} \rangle_{\text{th}}$ from [27–31] with a realistic estimate of the systematic error [35]: $\langle \sigma_{\text{CC}} \rangle_{\text{th}} = (9.2 \pm 0.5) \times 10^{-42} \text{ cm}^2$. The ν_e disappearance oscillation probability $P_{\nu_e \rightarrow \nu_x}$ can be written in terms of cross sections as

$$P_{\nu_e \rightarrow \nu_x} = 1 - \frac{\langle \sigma_{\text{CC}} \rangle_{\text{exp}}}{\langle \sigma_{\text{CC}} \rangle_{\text{th}}} \quad (5)$$

$$= 1 - \frac{9.4 \pm 0.89}{9.2 \pm 0.5} \quad (6)$$

where the statistical and systematical errors of the experimental value have been added quadratically. To get a density distribution for values of $P_{\nu_e \rightarrow \nu_x}$ we sampled $\langle \sigma_{\text{CC}} \rangle_{\text{exp}}$ and $\langle \sigma_{\text{CC}} \rangle_{\text{th}}$ from Gaussian distributions with the given mean values and widths (Fig. 6). The resulting distribution of $P_{\nu_e \rightarrow \nu_x}$ is slightly non Gaussian due to the ratio introduced in (6). Only positive values of $P_{\nu_e \rightarrow \nu_x}$ correspond to the physically allowed region of oscillation. To extract a 90% *CL* upper limit we therefore renormalize the distribution in $P_{\nu_e \rightarrow \nu_x}$ following the most conservative Bayesian approach [13]. The limit deduced is then

$$P_{\nu_e \rightarrow \nu_x} < 0.169 \quad (90\% \text{ CL}) \quad (7)$$

The interpretation of this limit in terms of the oscillation parameters Δm^2 and mixing angle is given in section IV.

B. Ratio of Cross Sections CC/NC

KARMEN measures, simultaneously, the CC reaction $^{12}\text{C}(\nu_e, e^-)^{12}\text{N}_{\text{g.s.}}$ with its sequential e^-/e^+ signature and the NC reaction $^{12}\text{C}(\nu, \nu')^{12}\text{C}^*(1^+, 1; 15.1 \text{ MeV})$ induced by $\nu_e, \bar{\nu}_\mu$ from μ^+ DAR. The NC event signature consists

¹The detection of $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillations by $p(\bar{\nu}_e, e^+)n$ leads to a different signature as the neutron capture time $\tau_n = 120 \mu\text{s}$ is much faster than the β -decays of ^{12}N or ^{12}B (0.5–36 ms time interval).

of a 15.1 MeV γ from deexcitation of $^{12}\text{C}^*$ detected within the $\nu_e, \bar{\nu}_\mu$ time window after beam-on-target. This reaction was observed by KARMEN for the first time [36] and is described in detail in [37,38]. The updated cross section for the 1990–1995 data

$$\langle \sigma_{\text{NC}} \rangle_{\text{exp}} = (10.8 \pm 0.9_{(\text{stat.})} \pm 0.8_{(\text{syst.})}) \times 10^{-42} \text{cm}^2 \quad (8)$$

again is in very good agreement with theoretical calculations (table I).

In the case of oscillations between ν_e and other ν flavors the NC cross section remains unchanged due to the flavor universality of NC interactions.² The number of CC events $^{12}\text{C}(\nu_e, e^-) ^{12}\text{N}_{\text{g.s.}}$ would decrease due to $\nu_e \rightarrow \nu_\tau$. By looking at the ratio of observed events (i.e. the measured cross sections) the systematic uncertainties in $\langle \sigma \rangle_{\text{exp}}$ are significantly reduced so that statistical fluctuations dominate the error on R_{exp} :

$$R_{\text{exp}} = \frac{\langle \sigma_{\text{CC}} \rangle_{\text{exp}}}{\langle \sigma_{\text{NC}} \rangle_{\text{exp}}} = 0.86 \pm 0.08 \quad (9)$$

This ν flux independent ratio R_{exp} is in good agreement with theoretical predictions (see table I). Again, there is no indication of ν_e disappearance.

As theoretical prediction for R_{th} we combine the R -values from [27–31] in a straight forward way and take the mean value and its variance as an estimate for the systematic error: $R_{\text{th}} = 0.91 \pm 0.035$. The $\nu_e \rightarrow \nu_x$ oscillation probability can then be written as

$$P_{\nu_e \rightarrow \nu_x} = 1 - R_{\text{exp}}/R_{\text{th}} \quad (10)$$

$$= 1 - \frac{0.86 \pm 0.08}{0.91 \pm 0.035} \quad (11)$$

In close analogy to the evaluation in section III A an upper limit can be extracted after renormalization to the physical region:

$$P_{\nu_e \rightarrow \nu_x} < 0.192 \quad (90\% \text{ CL}) \quad (12)$$

This limit is slightly higher than the limit (7) in section III A due to the relatively high value of the experimental cross section $\langle \sigma_{\text{NC}} \rangle_{\text{exp}}$ compared to the theoretical predictions.

C. Spectral Shapes

The third means of searching for oscillations $\nu_e \rightarrow \nu_x$ is to carry out a maximum likelihood (ML) analysis of the measured energy and spatial distribution of the prompt electrons from $^{12}\text{C}(\nu_e, e^-) ^{12}\text{N}_{\text{g.s.}}$ reactions. This method is not sensitive to the absolute number of detected sequences but to distortions of the expected spectra in energy and position within the detector volume. In order to minimize contributions from cosmic induced background we applied more stringent cuts in time (corresponding to $3\tau_{\mu^+}$) and spatial distribution (restricted fiducial volume of 86% of the central detector) for the prompt event resulting in 458 accepted sequences (see Fig. 7) with 5.8 background events determined in the appropriate prebeam evaluation. With a neutrino signal-to-background ratio of about 80:1 the cosmic background can be included as a fixed component in the likelihood analysis and will not be discussed further. The aim of the ML analysis is to calculate a combined likelihood for all 458 events on the basis that the two-dimensional density distribution $f(L, E_{e^-})$ is sensitive to varying amounts of oscillation events from $\nu_e \rightarrow \nu_\tau$, $\nu_e \rightarrow \nu_\mu$ and $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$. E_{e^-} is related to neutrino energy by $E_{e^-} = E_{\nu_e} - 17.3 \text{ MeV}$, $L_{e^-} = L_{\nu_e}$ since the electron is detected at the interaction point. The absolute energy scale is known to a precision of $\pm 0.25 \text{ MeV}$ from analysis of Michel electrons from decay of stopped cosmic muons. The density function f for no oscillations is

$$f_{\text{noosc.}}(L, E_{e^-}) = \frac{1}{F_{\text{noosc.}}} \cdot n(L) \cdot \Phi_{\nu_e}(E_{\nu_e}) \cdot \sigma(E_{\nu_e}) \quad (13)$$

with the normalization factor F integrated over the ν_e energy and the detector volume

²Note that this argument is only valid for flavor oscillations with neutrinos ν_e, ν_μ, ν_τ . Neutrino oscillations $\nu_e \rightarrow \nu_s$ involving sterile neutrinos would reduce both the number of ν_e -induced CC as well as $\nu_e, \bar{\nu}_\mu$ -induced NC reactions.

$$F_{noosc.} = \int_{E_{\nu_e}} \int_{Det} n(L) \cdot \Phi_{\nu_e}(E_{\nu_e}) \cdot \sigma(E_{\nu_e}) dL dE_{\nu_e} \quad (14)$$

The function $n(L)$ represents the spatial distribution of events within the rectangular detection volume resulting from an isotropic neutrino flux from ISIS ($\Phi_{\nu} \sim L^{-2}$). Φ_{ν_e} denotes the ν_e energy distribution from μ^+ DAR (see Fig. 1a), $\sigma(E_{\nu_e})$ the energy dependent cross section for $^{12}\text{C}(\nu_e, e^-)^{12}\text{N}_{g.s.}$ (see Fig. 4).

For an oscillation $\nu_e \rightarrow \nu_x$ with given Δm^2 the density function $f_{\Delta m^2}^{\nu}$ normalized by the integral $F_{\Delta m^2}^{\nu}$ over L and E can be written as

$$f_{\Delta m^2}^{\nu}(L, E_{e^-}) = \frac{1}{F_{\Delta m^2}^{\nu}} \cdot \sin^2(1.27 \frac{\Delta m^2 L}{E_{\nu_e}}) \cdot n(L) \cdot \Phi_{\nu_e}(E_{\nu_e}) \cdot \sigma(E_{\nu_e}) \quad (15)$$

The appearance of $\bar{\nu}_e$ through $\bar{\nu}_{\mu} \rightarrow \bar{\nu}_e$ is described by the density

$$f_{\Delta m^2}^{\bar{\nu}}(L, E_{e^+}) = \frac{1}{F_{\Delta m^2}^{\bar{\nu}}} \cdot \sin^2(1.27 \frac{\Delta m^2 L}{E_{\bar{\nu}_{\mu}}}) \cdot n(L) \cdot \Phi_{\bar{\nu}_{\mu}}(E_{\bar{\nu}_{\mu}}) \cdot \sigma(E_{\bar{\nu}_e}) \quad (16)$$

in close analogy to (15). The density functions used in the ML analysis are MC-simulated electron/positron (see Fig. 8) and positron/electron (see Fig. 3) distributions including all effects due to detector resolutions and the cuts applied. The combined likelihood for the $N = 458$ events is calculated with respect to the fraction r_1 of initially expected electrons now absent due to $\nu_e \rightarrow \nu_{\tau}$ and r_2 of the combined transition $\nu_e \rightarrow \nu_{\mu}$ and $\bar{\nu}_{\mu} \rightarrow \bar{\nu}_e$ taking into account the different detection efficiencies $\epsilon(\nu, \Delta m^2)$ and $\epsilon(\bar{\nu}, \Delta m^2)$ (see Fig. 5):

$$\begin{aligned} L_{\Delta m^2}(r_1, r_2) &= \prod_{k=1}^N L_{\Delta m^2}^k(r_1, r_2) = \prod_{k=1}^N \{ (1 + r_1 + r_2) \cdot f_{noosc.}(E_k, L_k) \\ &\quad - r_1 \cdot f_{\Delta m^2}^{\nu}(E_k, L_k) - r_2 \cdot \frac{1}{\epsilon(\nu, \Delta m^2) - \epsilon(\bar{\nu}, \Delta m^2)} \\ &\quad \cdot [\epsilon(\nu, \Delta m^2) \cdot f_{\Delta m^2}^{\nu}(E_k, L_k) - \epsilon(\bar{\nu}, \Delta m^2) \cdot f_{\Delta m^2}^{\bar{\nu}}(E_k, L_k)] \} \end{aligned} \quad (17)$$

In this specific construction of the likelihood function L the event number is fixed at $N = 458$, only the shape differences of the components are analyzed. Although not explicitly quoted in (17) we also include into $L_{\Delta m^2}(r_1, r_2)$ the knowledge, that the delayed energy E_{del} and the time difference t_{diff} differ for $^{12}\text{C}(\bar{\nu}_e, e^+)^{12}\text{B}$ (see Fig. 3).

Maximising $L_{\Delta m^2}(r_1, r_2)$ gives the most likely oscillation fractions $r_{1max}(\Delta m^2)$, $r_{2max}(\Delta m^2)$. This optimization is done separately for each value of Δm^2 in the range 0.1 to 100 eV²/c⁴. Over this range of Δm^2 the fully 2-dimensional best fit values are compatible with $r_1 = r_2 = 0$ within the 1 σ error band (see Fig. 9). As there is no indication for oscillations, upper limits $r_1(90\% CL)$, $r_2(90\% CL)$ for each Δm^2 are determined. In the case of $\nu_e \leftrightarrow \nu_{\tau}$ the ML analysis is not able to discriminate oscillation events for $2 \lesssim \Delta m^2 \lesssim 3 \text{ eV}^2/\text{c}^4$ and $\Delta m^2 > 50 \text{ eV}^2/\text{c}^4$. This is due to the fact that energy and spatial distributions for these oscillation cases are almost identical with the expected electron spectra without oscillations. However, for $\nu_e \leftrightarrow \nu_{\mu}$ mixing, the shape analysis is also sensitive for large Δm^2 since we expect events with $E_{pr} > 36 \text{ MeV}$ from $\bar{\nu}_{\mu} \rightarrow \bar{\nu}_e$.³ The 90% CL limits for each Δm^2 are calculated by integrating the corresponding 2-dimensional likelihood function for $r_1(r_2)$ over the whole $r_2(r_1)$ range and then renormalizing to the physical region $r_1(r_2) > 0$ applying a Bayesian approach near the physical boundary $r_1(r_2) = 0$, respectively. The limits obtained are equivalent to the limits $P_{\nu_e \rightarrow \nu_{\tau}}$ (90% CL), $P_{\nu_e \rightarrow \nu_{\mu}}$ (90% CL) in oscillation probability. The translation of these limits into 90% CL limits for the neutrino oscillation amplitude and mixing angles for different values of Δm^2 is shown in section IV.

IV. EXCLUSION LIMITS

In this section, we translate the results of the different methods obtained in section III as limits on the oscillation probability (cross section evaluation) or on the ν -mixings (shape analysis) into limits of the oscillation amplitude. This oscillation amplitude A is then expressed in terms of the mixing angles, for example $A^{e\tau} = \sin^2(2\Theta)$ (2 flavors) or $A^{e\tau} = 4\sin^2\Phi\cos^2\Phi\cos^2\Psi$ (3 flavors, see section I).

³The energy window $36 < E_{pr} < 50 \text{ MeV}$ can in fact be used to search for $\bar{\nu}_{\mu} \rightarrow \bar{\nu}_e$ oscillations in the appearance mode with $\bar{\nu}_e$ detection via $^{12}\text{C}(\bar{\nu}_e, e^+)^{12}\text{B}$.

A. Exclusion limits from measurement of the CC cross section

We first discuss the limit obtained from the evaluation of the absolute number of detected CC sequences (section III A): $P_{\nu_e \rightarrow \nu_x} < 0.169$ (90% CL)⁴ in a 2 flavor as well as in a complete 3 flavor scheme.

For a fixed neutrino energy and a fixed source–target distance L , the oscillation probability P is related to the mixing amplitude A via the spatial evolution term in (2). For an experimental configuration with continuous ν energies and a large volume detector with its resolution functions, this spatial evolution term is determined by MC simulations resulting in oscillation detection efficiencies $\epsilon(\nu, \Delta m^2)$ and $\epsilon(\bar{\nu}, \Delta m^2)$ as shown in Fig. 5. The upper limit of the oscillation amplitude for a given Δm^2 is then

$$A^{e\tau} \cdot \epsilon(\nu, \Delta m^2) < P_{\nu_e \rightarrow \nu_\tau} \text{ (90\% } CL) \quad (18)$$

For $\nu_e \rightarrow \nu_\mu$ the analogous MC simulations of positrons from $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ based on $f_{\Delta m^2}^{\bar{\nu}}$ were performed resulting in $\epsilon(\bar{\nu}, \Delta m^2)$ so that

$$A^{e\mu} \cdot (\epsilon(\nu, \Delta m^2) - \epsilon(\bar{\nu}, \Delta m^2)) < P_{\nu_e \rightarrow \nu_\mu} \text{ (90\% } CL) \quad (19)$$

$P_{\nu_e \rightarrow \nu_\tau}$ (90% CL) and $P_{\nu_e \rightarrow \nu_\mu}$ (90% CL) are the evaluated limits of the oscillation probability from the different analyses of section III.

Trivially, $P_{\nu_e \rightarrow \nu_x} = P_{\nu_e \rightarrow \nu_\tau} + P_{\nu_e \rightarrow \nu_\mu}$ assuming three neutrino flavors. However, in a 2 flavor scheme ignoring a second mixture, $P_{\nu_e \rightarrow \nu_x} = P_{\nu_e \rightarrow \nu_\tau}$ or $P_{\nu_e \rightarrow \nu_x} = P_{\nu_e \rightarrow \nu_\mu}$ respectively. Therefore the 90% CL limits in $\sin^2(2\Theta)$ can be evaluated following (18) and (19): $\sin^2(2\Theta) < 0.169/\epsilon(\nu, \Delta m^2)$ for $\nu_e \rightarrow \nu_\tau$ or $\sin^2(2\Theta) < 0.169/(\epsilon(\nu, \Delta m^2) - \epsilon(\bar{\nu}, \Delta m^2))$ for $\nu_e \rightarrow \nu_\mu$. Figure 10a shows the exclusion curves from this evaluation. Note that logical consistency requires that either one or the other limit holds, not both simultaneously. Parameter combinations of Δm^2 and $\sin^2(2\Theta)$ to the right of the curves are excluded at 90% CL . Taking $\nu_e \leftrightarrow \nu_\tau$ mixing, this corresponds to mixing angles of $\sin^2(2\Theta) > 0.338$ excluded for large Δm^2 and $\Delta m^2 > 0.77 \text{ eV}^2/\text{c}^4$ excluded for maximal mixing. Due to the very small efficiency $\epsilon(\nu, \Delta m^2) - \epsilon(\bar{\nu}, \Delta m^2)$ only three minor parameter regions can be excluded assuming $\nu_e \leftrightarrow \nu_\mu$ mixing. Figure 10b shows the KARMEN exclusion (curve 1) for $\nu_e \rightarrow \nu_\tau$ together with exclusion limits from other experiments. Curve 2 represents the 90% exclusion limit from a negative oscillation search in the channel $\bar{\nu}_e \rightarrow x$ at the nuclear power reactor of Bugey, France [15]. The second limit of a disappearance search comes from the GALLEX detector [16] in which the ν_e flux from the two ^{51}Cr source tests has been investigated⁵ with respect to $\nu_e \rightarrow \nu_x$. The Fermilab experiment E531 searched for CC τ -production through $\nu_e \rightarrow \nu_\mu$ and $\nu_e \rightarrow \nu_\tau$ appearance in a wideband $\nu_\mu(\nu_e)$ beam [18]. The exclusion limit from the $\nu_e \rightarrow \nu_\tau$ search is shown as curve 4. Due to the small experimental values $L/E < 0.1$ with a most probable value of $L/E \approx 0.01$, E531 is only sensitive to values of $\Delta m^2 > 10 \text{ eV}^2/\text{c}^4$.

In a more detailed 3 flavor evaluation, the KARMEN oscillation limit $P_{\nu_e \rightarrow \nu_x}$ corresponds to

$$A^{e\tau} \epsilon(\nu, \Delta m^2) + A^{e\mu} (\epsilon(\nu, \Delta m^2) - \epsilon(\bar{\nu}, \Delta m^2)) = P_{\nu_e \rightarrow \nu_x} < P_{\nu_e \rightarrow \nu_x} \text{ (90\% } CL) = 0.169 \quad (20)$$

In Fig. 11 the correlation of $A^{e\tau}$ and $A^{e\mu}$ limits is shown for some examples of Δm^2 . The parameter $A^{e\mu}$ is constrained only for regions at $\Delta m^2 \approx 2\text{--}3 \text{ eV}^2/\text{c}^4$, $\Delta m^2 \approx 7\text{--}9 \text{ eV}^2/\text{c}^4$ and at $\Delta m^2 \approx 13 \text{ eV}^2/\text{c}^4$ (see Fig. 10a), whereas for $\Delta m^2 \approx 5 \text{ eV}^2/\text{c}^4$ the upper limit of $A^{e\tau}$ even increases with $A^{e\mu}$ due to the fact that $\epsilon(\nu, \Delta m^2) - \epsilon(\bar{\nu}, \Delta m^2)$ is negative.

From $A^{e\tau} = 4 \sin^2 \Phi \cos^2 \Phi \cos^2 \Psi$ we can evaluate the 90% CL limits for the mixing angles Φ and Ψ taking the largest upper bound of $A^{e\tau}$ with respect to $A^{e\mu}$. With the double logarithmic presentation of $\tan^2 \Phi, \tan^2 \Psi$ we follow again [3]. Figure 12a,c shows the 90% CL limits in $A^{e\tau}$ and $A^{e\mu}$ from the cross section analysis as dashed curves (labeled with σ). In Fig. 12b, these limits from $A^{e\tau}$ are plotted as dashed curves for two examples of Δm^2 ($2 \text{ eV}^2/\text{c}^4$ and $100 \text{ eV}^2/\text{c}^4$) excluding areas in (Φ, Ψ) to the left. The corresponding limit on (Φ, Ψ) from $A^{e\mu}$ for $\Delta m^2 = 2 \text{ eV}^2/\text{c}^4$ is shown in Fig. 12d (dashed curve labeled σ).

The upper limit for $\nu_e \rightarrow \nu_\tau$, $P_{\nu_e \rightarrow \nu_\tau} < 0.169$ (90% CL), is reliable due to the spectroscopic measurement of the ν_e -flux from μ^+ DAR at ISIS and the nearly background-free $^{12}\text{C}(\nu_e, e^-)^{12}\text{N}_{\text{g.s.}}$ detection reaction. This limit is in contradiction to a model of 3 flavor neutrino mixing and masses calculated by Conforto [39,40]. Although being in

⁴Here, we do not consider the limit (12) from section IIIB since it is less stringent and therefore included in the limit used above.

⁵The complete analysis of the two ^{51}Cr source tests gives a ν signal of $R = 0.93 \pm 0.08$ compared with the expectation and results in slightly higher upper limits than an analysis [17] of a preliminary result of the first source test with $R = 1.04 \pm 0.12$.

conflict on a three standard deviations level with the negative result in $\nu_e \rightarrow \nu_\tau$ of FNAL E531 [18] this compilation of different experimental results predicts a transition probability $P_{\nu_e \rightarrow \nu_\tau} = 0.23 \pm 0.06$ (90% CL) expected in the appropriate ‘short-baseline’ regime. Figure 12a shows the calculated point $(\Delta m^2, A^{e\tau}) = (377 \pm 29 \text{ eV}^2/\text{c}^4, 0.46 \pm 0.12)$ lying completely in the area excluded at 90% CL by the KARMEN analysis of the absolute number of CC events.

B. Exclusion limits from the spectral shape analysis

In the 2-dimensional complete shape analysis (section III C) the oscillation probabilities $P_{\nu_e \rightarrow \nu_\mu}$ and $P_{\nu_e \rightarrow \nu_\tau}$ themselves depend on Δm^2 . The limit in the mixing amplitude is then

$$A^{e\tau} < \frac{r_1(90\% CL, \Delta m^2)}{\epsilon(\nu, \Delta m^2)} \quad \text{and} \quad A^{e\mu} < \frac{r_2(90\% CL, \Delta m^2)}{\epsilon(\nu, \Delta m^2) - \epsilon(\bar{\nu}, \Delta m^2)} \quad (21)$$

The resulting limits in $A^{e\tau}$ and $A^{e\mu}$ are shown in Figure 12a and Figure 12c, respectively, as solid exclusion curves (labeled ML) together with the exclusion curves evaluated above (dashed lines). For $\nu_e \rightarrow \nu_\tau$ the analysis of the spectral shape of L and E is only sensitive and competitive to the evaluation of the absolute event number in regions of about $0.5 \text{ E/L} < \Delta m^2 < 5 \text{ E/L}$ corresponding to Δm^2 values of ≈ 3 to $30 \text{ eV}^2/\text{c}^4$ for this experiment whereas for $\nu_e \leftrightarrow \nu_\mu$ the shape analysis is far more sensitive. This is due to expected events from $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ with positron energies $E_{e^+} > 36 \text{ MeV}$ (see also Fig. 3), where electrons from $^{12}\text{C}(\nu_e, e^-)^{12}\text{N}_{\text{g.s.}}$ cannot arise. Fig. 12b and Fig. 12d show the regions in (Φ, Ψ) space for various Δm^2 values and $\Delta m^2 = 2 \text{ eV}^2/\text{c}^4$, respectively, excluded by measurement of $\nu_e \leftrightarrow \nu_\tau$ and $\nu_e \leftrightarrow \nu_\mu$ mixing by KARMEN. These limits are compared with the exclusion limits from the Bugey $\bar{\nu}_e$ disappearance experiment where the particular value of Δm^2 chosen for comparison, $2 \text{ eV}^2/\text{c}^4$, has no special significance. Note that there are no limits on $A^{e\tau}$ from accelerator experiments like FNAL E531 below $\Delta m^2 = 10 \text{ eV}^2/\text{c}^4$.

C. Global limits from the 3 flavor analysis

Up to now, there are no bounds from $\nu_e \rightarrow \nu_\tau$ to the mixing angles Φ, Ψ in the region $\Delta m^2 \leq 10 \text{ eV}^2/\text{c}^4$ (see for example FNAL E531 limit in Fig. 10b). With $A^{e\tau} = 4U_{e3}^2 U_{\tau 3}^2 = 4\sin^2\Phi \cos^2\Phi \cos^2\Psi$ (see equ. 2) we can set new limits to the mixing angles Φ and Ψ from $\nu_e \rightarrow \nu_\tau$ oscillation search (see Fig. 12b, 13). For a given Δm^2 of for example $\Delta m^2 = 2 \text{ eV}^2/\text{c}^4$ the limits on the oscillation amplitudes $A^{e\tau}$ and $A^{e\mu}$ from the $\nu_e \rightarrow \nu_x$ search can be combined with other limits from KARMEN as well as from other experiments. Figure 13 demonstrates the complementarity of different experimental results with respect to the mixing angles.

The shaded area represents the 90% likelihood regions obtained by LSND [41] in favour of $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillations. The other curves show 90% CL limits from KARMEN ($\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ appearance search, $\nu_e \rightarrow \nu_\mu$ and $\nu_e \rightarrow \nu_\tau$ from this ν_e disappearance search), BNL E776 ($\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ appearance [42]) and FNAL E531 ($\nu_\mu \rightarrow \nu_\tau$ appearance [18]). For this value of Δm^2 , the $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ exclusion curve from BNL E776 just covers the allowed region by LSND⁶ with the KARMEN $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ sensitivity now substantially improved following an experimental upgrade (see section V). However, the complementary searches of KARMEN $\nu_e \rightarrow \nu_\tau$ and FNAL E531 $\nu_\mu \rightarrow \nu_\tau$ clearly exclude regions of the LSND evidence with $\tan^2\Phi \approx 1$ and $\tan^2\Psi \approx 1$, respectively. Again, the necessity of investigating all oscillation channels to aim for a consistent description of neutrino mixing becomes evident.

V. CONCLUSION AND OUTLOOK

The KARMEN experiment has found no positive evidence for ν oscillations in the investigated channels $\nu_e \rightarrow \nu_\tau$ and $\nu_e \rightarrow \nu_\mu$ through disappearance of ν_e . Charged current reactions induced by ν_e were analyzed by three different methods investigating the absolute number of events, the ratio of CC to NC events and the spectral shape of energy E and target distance L . The sensitivity of the $\nu_e \rightarrow \nu_x$ search is limited by the accuracy of the theoretical cross section of $^{12}\text{C}(\nu_e, e^-)^{12}\text{N}_{\text{g.s.}}$ and the ν flux calculation in the first evaluation, and by the relatively small statistics of 458 events in the case of the shape analysis. The experimental parameter L/E can be measured with high accuracy on an event

⁶Note that for values $\Delta m^2 < 2 \text{ eV}^2/\text{c}^4$ BNL E776 with its 90% CL limit does not cover the whole LSND region in (Φ, Ψ) any more.

basis and differs significantly compared to other $\bar{\nu}_e$ disappearance searches at reactors or $\nu_e \rightarrow \nu_\tau$ appearance searches at accelerators. In the maximum likelihood analysis, the 3 flavor approach assures fully independent treatment of $\nu_e \leftrightarrow \nu_\tau$ and $\nu_e \leftrightarrow \nu_\mu$ mixing as free parameters. The sensitivity of KARMEN is competitive with existing limits from oscillation searches at reactors and improves limits from accelerator experiments for $\nu_e \rightarrow \nu_\tau$ for $\Delta m^2 < 10 \text{ eV}^2/\text{c}^4$.

In 1996, the KARMEN detector system was improved by an additional veto counter system consisting of 300 m^2 of plastic scintillator modules placed within the massive iron shielding of the central detector. This new veto system will reduce significantly cosmic induced background for the oscillation search, especially in the appearance channel $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ [43]. In the ongoing measuring period we expect again about 400 ν_e -induced CC sequences from $^{12}\text{C}(\nu_e, e^-)^{12}\text{N}_{\text{g.s.}}$. Therefore, the statistical significance of this $\nu_e \rightarrow \nu_x$ investigation will slightly improve ending up with about 900 ν_e -induced CC reactions in the KARMEN detector after a measuring period of 2–3 years.

We gratefully acknowledge the financial support of the German Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie (BMBF), the Particle and Astronomy Research Council (PPARC) and the Central Laboratory of the Research Council (CLRC) in the UK.

* Present Address: Department of Physics and Astronomy, University of Alabama, Tuscaloosa AL 35487, USA

- [1] A. Acker and S. Pakvasa, Phys. Lett. **B 397** (1997) 209
- [2] P.F. Harrison et al., Phys. Lett. **B 349** (1995) 137
Phys. Lett. **B 396** (1997) 186
- [3] G. Fogli et al., Phys. Rev. **D 52** (1995) 5334
Phys. Rev. **D 56** (1997) 3081
- [4] S.M. Bilenky et al., Phys. Lett. **B 356** (1995) 273
Phys. Rev. **D 54** (1996) 1881
- [5] K.S. Babu et al., Phys. Lett. **B 359** (1995) 351
- [6] E. Torrente-Lujan, Phys. Lett. **B 389** (1996) 557
- [7] H. Minakata, Phys. Rev. **D 52** (1995) 6630
- [8] S. Barshay and P. Heiliger, Astropart. Phys. **6** (1997) 323
- [9] C.Y. Cardall and G.M. Fuller, Phys. Rev. **D 53** (1996) 4421
- [10] D. Suematsu, Phys. Lett. **B 392** (1997) 413
- [11] Z.G. Bereshiani and R.N. Mohapatra, Phys. Rev. **D 52** (1995) 6607
- [12] R. Foot and R.R. Volkas, Phys. Rev. **D 52** (1995) 6595
- [13] Particle Data Group, Phys. Rev. **D 54** (1996) 94
- [14] B. Zeitnitz et al., to be published in Prog. Part. Nucl. Physics **40** (1997)
- [15] B. Achkar et al., Nucl. Phys. **B 434** (1995) 503
- [16] W. Hampel et al. (GALLEX Collaboration), MPI H-V41–1997, submitted to Phys. Lett. **B**
- [17] J. Bahcall et al., Phys. Lett. **B 348** (1995) 121
- [18] N. Ushida et al., Phys. Rev. Lett. **57** (1986) 2897
- [19] R.L. Burman et al., Nucl. Instr. Meth. **A 368** (1996) 416
- [20] C. Bouchiat et al., Phys. Rev. **106** (1957) 170
- [21] G. Drexlin et al., Nucl. Instr. Meth. **A 289** (1990) 490
- [22] A. Dodd et al., Nucl. Instr. Meth. **A 336** (1993) 136
- [23] K. Eitel et al., Nucl. Instr. Meth. **A 340** (1994) 346
- [24] B. Bodmann et. al., Phys. Lett. **B 280** (1992) 198
- [25] B.A. Bodmann et. al., Phys. Lett. **B 339** (1994) 215
- [26] B. Armbruster et. al., to be submitted to Phys. Lett. **B**
- [27] M. Fukugita et al., Phys. Lett. **B 212** (1988) 139
- [28] S. Mintz and M. Pourkaviani, Phys. Rev. **C 40** (1989) 2458
- [29] E. Kolbe et al., Phys. Rev. **C 49** (1994) 1122
- [30] J. Engel et al., Phys. Rev. **C 54** (1996) 2740
- [31] J.S. O’Connell et al., Phys. Rev. **C 6** (1972) 719
- [32] T.W. Donnelly and R. Peccei, Phys. Rep. **50** (1979) 1
T.W. Donnelly, private communication, 1991
- [33] J. Bernab   and P. Pascual, Nucl. Phys. **A 324** (1979) 365
- [34] R. Parthasarathy, Int. J. Mod. Phys. **E 3** (1994) 1195
- [35] P. Vogel, private communication (1996)

- [36] B. Bodmann et. al., Phys. Lett. **B 267** (1991) 321
[37] B.E. Bodmann et. al., Phys. Lett. **B 332** (1994) 251
[38] R. Maschuw et al., to be published in Prog. Part. Nucl. Physics, **40** (1997)
[39] G. Conforto, Il Nuovo Cimento **103 A** (1990) 751
[40] G. Conforto et al., Astropart. Phys. **5** (1996) 147
[41] C. Athanassopoulos et al., Phys. Rev. **C 54** (1996) 2685
[42] L. Borodovsky et al., Phys. Rev. Lett. **68** (1992) 274
[43] G. Drexlin et al., to be published in Prog. Part. Nucl. Physics **40** (1997)

model	$^{12}\text{C}(\nu_e, e^-)^{12}\text{N}_{\text{g.s.}}$	$^{12}\text{C}(\nu, \nu')^{12}\text{C}^*$	R	reference
EPT ⁷	9.19	9.87	0.93	Fukugita [27]
EPT	9.0	10.6	0.85	Mintz [28]
CRPA ⁸	9.29	10.53	0.88	Kolbe [29]
Shell with OBD ⁹	9.1	9.8	0.93	Engel [30]
Shell with OBD ¹⁰	9.4	9.9	0.95	O'Connell [31]
EPT		10.3		Bernabéu [33]
Shell		9.9		Parthasarathy [34]
	$9.4 \pm 0.4 \pm 0.8$	$10.8 \pm 0.9 \pm 0.8$	0.86 ± 0.08	this experiment

TABLE I. Comparison of theoretical cross sections $\langle \sigma \rangle_{\text{th}}$ in units of 10^{-42}cm^2 averaged over the neutrino energy from μ^+ DAR and the values $\langle \sigma \rangle_{\text{exp}}$ measured with KARMEN (including statistical and systematical errors); for the NC cross section $\nu = \nu_e + \bar{\nu}_\mu$ in case of no oscillations; R indicates the ratio $\sigma_{CC}(\nu_e)/\sigma_{NC}(\nu_e + \bar{\nu}_\mu)$.




FIG. 1. Neutrino energy spectra (a) and production time (b) at ISIS. The proton double pulses are repeated with a frequency of 50 Hz.

⁷Elementary Particle Treatment

⁸Continuum Random Phase Approximation

⁹One Body Density

¹⁰This value for the CC exclusive cross section is calculated by [30] using the OBD of [31], but is also obtained with the original computer code NUCS from [32].




FIG. 2. Signatures of the ν_e detection reaction $^{12}\text{C}(\nu_e, e^-)^{12}\text{N}_{\text{g.s.}}$: time and energy distributions of the prompt electron (a;c) and the delayed positron (b;d). Data points are with background subtracted, MC expectations are shown as histograms, total background as shaded histograms.




FIG. 3. Simulated detector response in energy, target distance and time difference including readout efficiency of $^{12}\text{C}(\nu_e, e^-)^{12}\text{N}_{\text{g.s.}}$ (solid histograms) and $^{12}\text{C}(\bar{\nu}_e, e^+)^{12}\text{B}$ (dashed histograms) for $\Delta m^2 = 100 \text{ eV}^2/c^4$ normalized to the same arbitrary number of events.




FIG. 4. Energy dependence of the cross sections of charged current (CC) reactions $^{12}\text{C}(\nu_e, e^-)^{12}\text{N}_{\text{g.s.}}$ and $^{12}\text{C}(\bar{\nu}_e, e^+)^{12}\text{B}$ and of neutral current (NC) $^{12}\text{C}(\nu, \nu')^{12}\text{C}^*(1^+, 1; 15.1 \text{ MeV})$ for different ν flavors in an elementary particle treatment of the Carbon nucleus [27].




FIG. 5. Relative detection efficiencies ϵ as functions of the mass parameter Δm^2 : a) $\epsilon(\nu)$ for $\nu_e \leftrightarrow \nu_\tau$ mixing; b) $\nu_e \leftrightarrow \nu_\mu$ mixing; curve 1 shows $\epsilon(\nu)$ for $\nu_e \rightarrow \nu_\mu$ disappearance, curve 2 shows $\epsilon(\bar{\nu})$ for $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ appearance; curve 1-2 shows the net efficiency $\epsilon(\nu) - \epsilon(\bar{\nu})$ for the combination of $\nu_e \rightarrow \nu_\mu$ and $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$. The efficiencies ϵ are normalized to the detection efficiency of $^{12}\text{C}(\nu_e, e^-)^{12}\text{N}_{\text{g.s.}}$ sequences within the applied cuts.




FIG. 6. Simulated distributions of cross sections $\langle \sigma_{\text{CC}} \rangle_{\text{exp}}$ and $\langle \sigma_{\text{CC}} \rangle_{\text{th}}$ of $^{12}\text{C}(\nu_e, e^-)^{12}\text{N}_{\text{g.s.}}$ and resulting $P_{\nu_e \rightarrow \nu_x}$ according to Gaussian distributed values of $\langle \sigma_{\text{CC}} \rangle$. The shaded areas show the physical region, darkly shaded are the upmost 10% of the shaded $P_{\nu_e \rightarrow \nu_x}$ distribution.




FIG. 7. Distributions of energy and target distance of prompt events (data points) with stringent cuts for the $\nu_e \rightarrow \nu_x$ shape analysis and underlying MC expectations without oscillations.




FIG. 8. Examples of simulated e^- energy and distance distributions (E - and L -projection of $f_{noosc.} - f_{\Delta m^2}^\nu$) in the oscillation scenario $\nu_e \rightarrow \nu_\tau$ for maximal mixing $A^{e\tau} = 1$ with different Δm^2 as well as for no oscillation. The spectra show the visible electron signal in the detector and are normalized to the experimental event number $N = 458$ for no oscillation (see Fig. 7).




FIG. 9. Result of the 2-dimensional ML shape analysis of the 458 events in the channels $\nu_e \rightarrow \nu_\tau$ and $\nu_e \rightarrow \nu_\mu + \bar{\nu}_\mu \rightarrow \bar{\nu}_e$. Shown are the best fit values for oscillation contributions (solid lines) with the corresponding $1\text{-}\sigma$ errors (dashed lines) as a function of the oscillation parameter Δm^2 .




FIG. 10. 2 flavor sensitivity plot for the different channels $\nu_e \rightarrow \nu_\mu$ and $\nu_e \rightarrow \nu_\tau$ from KARMEN (a) and comparison with limits of other experiments (b).

a: KARMEN limits from the absolute cross section for $\nu_e \leftrightarrow \nu_\tau$ (solid line) and $\nu_e \leftrightarrow \nu_\mu$ (dashed line) mixing.

b: Limits from KARMEN $\nu_e \rightarrow \nu_\tau$ (1), Bugey $\bar{\nu}_e \rightarrow \bar{\nu}_x$ (2), GALLEX $\nu_e \rightarrow \nu_x$ (3) and FNAL E531 $\nu_e \rightarrow \nu_\tau$ (4).




FIG. 11. 3 flavor 90% *CL* exclusion limits of the mixing amplitudes $A^{e\mu}$ and $A^{e\tau}$ based on the extracted oscillation limit of $P_{\nu_e \rightarrow \nu_x} < 0.169$ (90% *CL*) from the cross section analysis (see equ. 7). Areas to the right of the lines are excluded.




FIG. 12. a,c: KARMEN exclusion limits of the mixing amplitude $A^{e\tau}$ and $A^{e\mu}$ (solid line = shape ML; dashed line (σ) = absolute number of events)
b,d: corresponding limits of the mixing angles Ψ and Φ from the limits of: (b) $A^{e\tau}$ for several values of Δm^2 , (d) $A^{e\mu}$, for $\Delta m^2 = 2 \text{ eV}^2/\text{c}^4$. For comparison, the exclusion limits from $\bar{\nu}_e$ disappearance search at Bugey [15] are shown, also for $\Delta m^2 = 2 \text{ eV}^2/\text{c}^4$. In b(d) areas to the left(right) of the curves, respectively, and between the horizontal lines are excluded. In (d) curves 1 and 2 are the KARMEN limits from $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ and $\nu_\mu \rightarrow \nu_e$ appearance search.




FIG. 13. 90% CL limits to the mixing angles Φ and Ψ fixed at $\Delta m^2 = 2\text{eV}^2/c^4$ deduced from this experiment from the oscillation channels $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ appearance and $\nu_e \rightarrow \nu_\mu$, $\nu_e \rightarrow \nu_\tau$ disappearance in comparison with other experiments: FNAL E531 $\nu_\mu \rightarrow \nu_\tau$ appearance [18], BNL E776 $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ appearance [42] and a 90% likelihood favoured region from LSND [41] (shaded region).